

No. of Questions: 25

Answers: Available

Format: Pdf/Txt

Downloadable: Yes

Prepared by: Sathish

NVS staff nurse question paper Part-2 from Question No.26-50

Question No. *26. The commonest cause of bile duct strictures (a) Sclerosing cholangitis

(b) Post cholecystectomy

(c) Malignancy

(d) Ascending cholangitis

Question No. *27. Commonest indication of surgery in chronic pancreatitis?

(a) Pain

(b) Infection

(c) Malignant transformation

(d) Duodenal obstruction

Question No. *28. Congenital atresia is commonest in?

(a) Distal colon (b) Duodenum

(c) Proximal colon (d) Ileum

Question No. *29. Visceral aneurysm is most common in

(a) Renal artery (b) Hepatic artery

(c) Splenic artery (d) Left gastric artery

Question No. *30. The following drugs used in treatment of cyclical mastalgia except

- (a) Estrogen
- (b) Danazol
- (c) Tamoxifen
- (d) Evening primrose oil

Question No. *31. What is the earliest indication of volkmann's ischemia?

- (a) Pain
- (b) Pallor & Poor Capillary filling
- (c) Paraesthesia in median nerve area
- (d) Contracture of finger

Question No. *32. In osteomalacia

- (a) Osteoid formation is defective
- (b) Mineralisation of bone is defect
- (c) Both
- (d) None

Question No. *33. Rickets is characterized by

- (a) Decrease calcium content of bone
- (b) Widening of growth cartilage
- (c) Beading ribs
- (d) All of above

Question No. *34. The most radiosensitive tumor is

- (a) Ewing's Sarcoma
- (b) Osteosarcoma

(c) Fibrosarcoma (d) Liposarcoma

Question No. *35. Meniscal injury occurs during

(a) Rotation

(b) Extention

(c) Rotation & Flexion

(d) Flexion

Question No. *36. Fracture of first metacarpal bone

(a) Barton's fracture

(b) Bennett's fracture

(c) Chauffer's fracture

(d) Boxer's fracture

Question No. *37. Functions of ovary include all of the following except

(a) Autocrine (b) Paracrine

(c) Exocrine (d) Endocrine

Question No. *38. Home essential for the maintenance of pregnancy

(a) LH (b) Progestrone

(c) HCG (d) Estrogen

Question No. *39. Major function of von willebrand factor is

(a) Platelet adhesion

(b) Binding factor VIII

(c) Activation of factor XII

www.nursingwork.in

www.rightjobalert.com

(d) Stimulation of platelet release

Question No. * 40. Blue nevi and multiple lentiginos are associated with the following type of cardiac disorder

- (a) Cardiomyopathy
- (b) Atrial myxoma
- (c) Constrictive pericardity
- (d) Cardiac arrhythmias

Question No. *41. The prognosis of the newborn of a lady with accidental hemorrhage depends on

- (a) Admission delivery interval
- (b) Weight of the body
- (c) Abruptio delivery interval
- (d) Volume of retroplacental clots

Question No. *42. The treatment of choice for hydatidiform mole

- (a) Hysterotomy
- (b) Medical treatment
- (c) Hysterectomy
- (d) Suction evaluation

Question No. *43. Which scar is least prone for rupture

- (a) Classic scare
- (b) I shaped scar
- (c) LSCS

www.nursingwork.in

www.rightjobalert.com

- (d) Inverted T-shaped scar

Question No. *44. Subnuclear vacuolation of endometrium is seen in

- (a) Menstrual phase
- (b) Ovarian phase
- (c) Proliferatory phase
- (d) Secretory phase

Question No. *45. Ergometrine is contra indicated in all except

- (a) Eclampsia
- (b) Rheumatic heart disease
- (c) Moderate anemia
- (d) Rh incompatibility

Question No. *46. All of the following are true regarding asymmetric IUGR except

- (a) Brain is spared
- (b) PIH is a cause
- (c) Occur in later part of pregnancy
- (d) Chromosomal anomalies occur

Question No. *47. Monochorionic monoamniotic twins occur due to division during

- (a) Within 24 hours
- (b) 24-72 hours
- (c) 72 hours to 8 days
- (d) After 8 days

Question No. *48. Most common site of puerperal infection

- (a) Placental site

www.nursingwork.in

www.rightjobalert.com

- (b) Cervical laceration
- (c) Episiotomy wound
- (d) Vaginal laceration

Question No. *49. Acute cryptomenorrhea can occur due to all of the following except

- (a) Conization
- (b) Trachelorrhaphy

- (c) Cryosurgery
- (d) Amputation of cervix

Question No. *50. An example of type-I respiratory failure

- (a) ARDS
- (b) Chronic bronchitis
- (c) Narcotic overdose
- (d) Primary alveolar hypoventilation

Answers for Above Questions

- Answer for Question No. *26 Option B
- Answer for Question No. *27 Option A
- Answer for Question No. *28 Option B
- Answer for Question No. *29 Option C
- Answer for Question No. *30 Option A
- Answer for Question No. *31 Option A
- Answer for Question No. *32 Option B
- Answer for Question No. *33 Option D
- Answer for Question No. *34 Option A
- Answer for Question No. *35 Option C
- Answer for Question No. *36 Option B
- Answer for Question No. *37 Option C
- Answer for Question No. *38 Option B
- Answer for Question No. *39 Option A
- Answer for Question No. *40 Option B
- Answer for Question No. *41 Option A
- Answer for Question No. *42 Option D
- Answer for Question No. *43 Option C
- Answer for Question No. *44 Option D
- Answer for Question No. *45 Option C
- Answer for Question No. *46 Option D
- Answer for Question No. *48 Option A
- Answer for Question No. *47 Option D
- Answer for Question No. *49 Option C
- Answer for Question No. *50 Option A

www.nursingwork.in

www.rightjobalert.com